

(3)

Theorem [Eholzer - n]

Let C be integers, H_C set of cusp. conf. den.

Assume there exist f_h ($h \in H_C$) satisfying (1) - (5)

and such that f_h is inv. under some congruence subgroup of Γ .

Then the functions f_h are unique (up to multiplic. by scalar)

Three main ingredients in proof

1st

Let $\xi =$ column vector of the f_h (same fixed order), then

$$\xi(A\tau) = \rho(A) \xi(\tau) \text{ for a rep. } \rho: \Gamma \rightarrow GL(n, \mathbb{C}) \text{ (} n = \#H_C \text{)}$$

vector valued modular function on Γ

useful to multiply by powers of

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$$

then we land in a

space of vector valued modular forms on Γ (poss. of half-integral wt)

$$\tilde{F} = \{ (A, w(\tau)) \mid A \in \Gamma, w \text{ vector w.t. } \text{Hol}(\rho), w^2 = c\tau + d \}$$

$$\text{Group: } (A, w) \cdot (A', w') = (AA', w(A'\tau)w'(\tau))$$

$$\text{action: } F \in \mathfrak{g} \rightarrow \mathbb{C}^n, (A, w) \in \tilde{F}: F|_k(A, w) = F(A\tau) w(\tau)^{-2k}$$

$$\rho: \tilde{F} \rightarrow GL(n, \mathbb{C}) \text{ representation}$$

$$M_k(\rho) = \left\{ F: \mathfrak{g} \xrightarrow{\text{hol}} \mathbb{C}^n \mid F|_k \alpha = \rho(\alpha) F \forall \alpha \in \tilde{F} \right. \\ \left. F = O(1) \text{ for } \text{Im } z \rightarrow 0 \right\}$$

basic example:

$$\eta \in M_{1/2}(\theta) \text{ for a } \theta: \tilde{F} \rightarrow \mathbb{P}^1$$

$$\eta^k \in M_{k/2}(\theta^k \otimes \rho) \text{ for } k \in \mathbb{Z} \text{ } k \geq 0$$